

SUBJECT INDEX Volume 63

Actins

Etienne P., Parés-Herbuté N., Mani-Ponset L., Gabrion J., Rabesandratana H., Herbuté S., Monnier L.
Phenotype modulation in primary cultures of aortic smooth muscle cells from streptozotocin-diabetic rats, 63:223–234

Adenocarcinoma cells

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.
Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305–317

Adipose tissue

Nakajima I., Yamaguchi T., Ozutsumi K., Aso H.
Adipose tissue extracellular matrix: newly organized by adipocytes during differentiation, 63:193–200

Alveolar formation

Schünemann H.J., Dillon D., Nielsen L.C., Lwebuga-Mukasa J.S.
Modulation of laminin integrin receptors in the postnatal and adult rat lung, 63:181–191

Ambystoma mexicanum

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.
Post-transcriptional control of *c-myc* RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69–79

Amphibian development

Ishizuya-Oka A., Inokuchi T., Ueda S.
Thyroid hormone-induced apoptosis of larval cells and differentiation of pepsinogen-producing cells in the stomach of *Xenopus laevis* in vitro, 63:59–68

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.
Post-transcriptional control of *c-myc* RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69–79

Ohmura H., Wakahara M.

Transformation of skin from larval to adult types in normally metamorphosing and metamorphosis-arrested salamander, *Hynobius retardatus*, 63:237–246

Mizuno N., Mochii M., Takagi C., Takahashi T.C., Eguchi G., Okada T.S.

A critical role for the optic vesicle in lens development: A reinvestigation of free lens formation in *Cynops pyrrhogaster*, 63:247–252

Antiproliferative effects

Vrana J.A., Saunders A.M., Chellappan S.P., Grant S.
Divergent effects of bryostatin 1 and phorbol myristate acetate on cell cycle arrest and maturation in human myelomonocytic leukemia cells (U937), 63:33–42

Apoptosis

Sreng L.
Apoptosis-inducing brain factors in maturation of an insect sex pheromone gland during differentiation, 63:53–58

Ishizuya-Oka A., Inokuchi T., Ueda S.
Thyroid hormone-induced apoptosis of larval cells and differentiation of pepsinogen-producing cells in the stomach of *Xenopus laevis* in vitro, 63:59–68

Ohmura H., Wakahara M.

Transformation of skin from larval to adult types in normally metamorphosing and metamorphosis-arrested salamander, *Hynobius retardatus*, 63:237–246

Atherosclerosis

Etienne P., Parés-Herbuté N., Mani-Ponset L., Gabrion J., Rabesandratana H., Herbuté S., Monnier L.
Phenotype modulation in primary cultures of aortic smooth muscle cells from streptozotocin-diabetic rats, 63:223–234

Axolotl

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.
Post-transcriptional control of *c-myc* RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69–79

Basonuclin

Weiner L., Green H.
Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263–272

Bone alkaline phosphatase

Siggelkow H., Niedhart C., Kurre W., Ihbe A., Schulz A., Atkinson M.J., Hüfner M.
In vitro differentiation potential of a new human osteosarcoma cell line (HOS 58), 63:81–91

Brain factors

Sreng L.
Apoptosis-inducing brain factors in maturation of an insect sex pheromone gland during differentiation, 63:53–58

Bryostatin

Vrana J.A., Saunders A.M., Chellappan S.P., Grant S.
Divergent effects of bryostatin 1 and phorbol myristate acetate on cell cycle arrest and maturation in human myelomonocytic leukemia cells (U937), 63:33–42

Bullous pemphigoid antigen, BP-230

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.
Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1–11

Cadherins

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.
Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1–11

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295–304

Cancer

Dumortier J., Daemi N., Pourreyron C., Anderson W., Bellaton C., Jacquier M.-F., Bertrand S., Chayvialle J.-A., Remy L.

Loss of epithelial differentiation markers and acquisition of vimentin expression after xenograft with laminin-1 enhance migratory and invasive abilities of human colon cancer cells LoVo C5, 63:141-150

Catenins

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.

Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Cattle

Nakajima I., Yamaguchi T., Ozutsumi K., Aso H.

Adipose tissue extracellular matrix: newly organized by adipocytes during differentiation, 63:193-200

CD 15

Nohr D., Mai J.K.

Distribution of the CD15 epitope in the mammalian developing lung is opposite in mouse compared with human, 63:43-49

Cell culture

Siggelkow H., Niedhart C., Kurre W., Ihbe A., Schulz A., Atkinson M.J., Hüfner M.

In vitro differentiation potential of a new human osteosarcoma cell line (HOS 58), 63:81-91

Cohen-Tanugi A., Forest N.

Retinoic acid suppresses the osteogenic differentiation capacity of murine osteoblast-like 3A/1D-1M cell cultures, 63:115-123

Stingl J., Eaves C.J., Kuusk U., Emerman J.T.

Phenotypic and functional characterization in vitro of a multipotent epithelial cell present in the normal adult human breast, 63:201-212

Etienne P., Parés-Herbuté N., Mani-Ponset L., Gabrion J., Rabesandratana H., Herbuté S., Monnier L.

Phenotype modulation in primary cultures of aortic smooth muscle cells from streptozotocin-diabetic rats, 63:223-234

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.

Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.

Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

Cell-matrix interaction

Schünemann H.J., Dillon D., Nielsen L.C., Lwebuga-Mukasa J.S. Modulation of laminin integrin receptors in the postnatal and adult rat lung, 63:181-191

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.

Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

c-myc

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.

Post-transcriptional control of c-myc RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69-79

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.

Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Cockroach

Sreng L.

Apoptosis-inducing brain factors in maturation of an insect sex pheromone gland during differentiation, 63:53-58

Collagens

Stingl J., Eaves C.J., Kuusk U., Emerman J.T.

Phenotypic and functional characterization in vitro of a multipotent epithelial cell present in the normal adult human breast, 63:201-212

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.

Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Fröjdman K., Pelliniemi L.J., Virtanen I.

Differential distribution of type IV collagen chains in the developing rat testis and ovary, 63:125-130

Colon

Dumortier J., Daemi N., Pourreyron C., Anderson W., Bellaton C., Jacquier M.-F., Bertrand S., Chayvialle J.-A., Remy L.

Loss of epithelial differentiation markers and acquisition of vimentin expression after xenograft with laminin-1 enhance migratory and invasive abilities of human colon cancer cells LoVo C5, 63:141-150

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.

Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

Crystallin genes

Mizuno N., Mochii M., Takagi C., Takahashi T.C., Eguchi G., Okada T.S.

A critical role for the optic vesicle in lens development; A reinvestigation of free lens formation in *Cynops pyrrhogaster*, 63:247-252

Cyclin D

Goumans M.-J., Ward-van Oostwaard D., Wianny F., Savatier P., Zwijnen A., Mummery C.

Mouse embryonic stem cells with aberrant transforming growth factor β signalling exhibit impaired differentiation in vitro and in vivo, 63:101-113

Cynops

Mizuno N., Mochii M., Takagi C., Takahashi T.C., Eguchi G., Okada T.S.

A critical role for the optic vesicle in lens development; A reinvestigation of free lens formation in *Cynops pyrrhogaster*, 63:247-252

Cytoskeleton

Fabre-Jonca N., Allaman J.-M., Radlgruber G., Meda P., Kiss J.Z., French L.E., Masson D.

The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Dedifferentiation

Cohen-Tanugi A., Forest N.

Retinoic acid suppresses the osteogenic differentiation capacity of murine osteoblast-like 3A/1D-1M cell cultures, 63:115-123

Etienne P., Parés-Herbuté N., Mani-Ponset L., Gabrion J., Rabesandratana H., Herbuté S., Monnier L.

Phenotype modulation in primary cultures of aortic smooth muscle cells from streptozotocin-diabetic rats, 63:223-234

Desmocollins

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Desmogleins

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Desmosomes

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.
Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Development

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.
Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.
Post-transcriptional control of *c-myc* RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69-79

Fröjdman K., Pelliniemi L.J., Virtanen I.
Differential distribution of type IV collagen chains in the developing rat testis and ovary, 63:125-130

Hayward S.W., Haughey P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.
Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Kloth S., Gmeiner T., Aigner J., Jennings M.L., Röckl W., Minuth W.W.
Transitional stages in the development of the rabbit renal collecting duct, 63:21-32

Diabetes

Etienne P., Parés-Herbuté N., Mani-Ponset L., Gabrion J., Rabesandratana H., Herbuté S., Monnier L.
Phenotype modulation in primary cultures of aortic smooth muscle cells from streptozotocin-diabetic rats, 63:223-234

Differentiation markers

Dumortier J., Daemi N., Pourreya C., Anderson W., Bellaton C., Jacquier M.-F., Bertrand S., Chayvialle J.-A., Remy L.
Loss of epithelial differentiation markers and acquisition of vimentin expression after xenograft with laminin-1 enhance migratory and invasive abilities of human colon cancer cells LoVo C5, 63:141-150

Embryonic stem cell

Goumans M.-J., Ward-van Oostwaard D., Wianny F., Savatier P., Zwijsen A., Mummery C.
Mouse embryonic stem cells with aberrant transforming growth factor β signalling exhibit impaired differentiation in vitro and in vivo, 63:101-113

Epidermal tumors

Magnaldo T., Fowles D., Darmon M.
Galectin-7, a marker of all types of stratified epithelia, 3:159-168

Epidermis

Magnaldo T., Fowles D., Darmon M.
Galectin-7, a marker of all types of stratified epithelia, 3:159-168

Weiner L., Green H.

Basoonin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Epithelial differentiation

Hayward S.W., Haughey P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.
Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Kloth S., Gmeiner T., Aigner J., Jennings M.L., Röckl W., Minuth W.W.
Transitional stages in the development of the rabbit renal collecting duct, 63:21-32

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.
Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

Magnaldo T., Fowles D., Darmon M.
Galectin-7, a marker of all types of stratified epithelia, 63:159-168

Fabre-Jonca N., Allaman J.-M., Radlgruber G., Meda P., Kiss J.Z., French L.E., Masson D.
The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.
Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Zvibel I., Fiorino A.S., Brill S., Reid L.M.
Phenotypic characterization of hepatoma cell lines and lineage-specific regulation of gene expression by differentiation agents, 63:213-221

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.
Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

Extracellular matrix

Nakajima I., Yamaguchi T., Ozutsumi K., Aso H.
Adipose tissue extracellular matrix: newly organized by adipocytes during differentiation, 63:193-200

Fibronectin

Galdieri M., Ricci G.
Characterization of different cell populations isolated from rat testis peritubular cells, 63:13-19

Flow cytometry

Stingl J., Eaves C.J., Kuusk U., Emerman J.T.
Phenotypic and functional characterization in vitro of a multipotent epithelial cell present in the normal adult human breast, 63:201-212

Focal adhesion kinase (FAK)

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C. Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

Free Lens

Mizuno N., Mochii M., Takagi C., Takahashi T.C., Eguchi G., Okada T.S.
A critical role for the optic vesicle in lens development; A reinvestigation of free lens formation in *Cynops pyrrhogaster*, 63:247-252

Galectin-7

Magnaldo T., Fowles D., Darmon M.
Galectin-7, a marker of all types of stratified epithelia, 63:159-168

Gene expression

Zvibel I., Fiorino A.S., Brill S., Reid L.M.
Phenotypic characterization of hepatoma cell lines and lineage-specific regulation of gene expression by differentiation agents, 63:213-221

Mizuno N., Mochii M., Takagi C., Takahashi T.C., Eguchi G., Okada T.S.
A critical role for the optic vesicle in lens development; A reinvestigation of free lens formation in *Cynops pyrrhogaster*, 63:247-252
Krause R., Hemberger M., Messerschmid M., Mayer W., Kothary R., Dixkens C., Fundele R.
Molecular cloning and characterization of murine *Mpgc60*, a gene predominantly expressed in the intestinal tract, 63:285-294

Gland differentiation

Sreng L.
Apoptosis-inducing brain factors in maturation of an insect sex pheromone gland during differentiation, 63:53-58

Goblet cells

Krause R., Hemberger M., Messerschmid M., Mayer W., Kothary R., Dixkens C., Fundele R.
Molecular cloning and characterization of murine *Mpgc60*, a gene predominantly expressed in the intestinal tract, 63:285-294

Growth factors

Goumans M.-J., Ward-van Oostwaard D., Wianny F., Savatier P., Zwijnen A., Mummery C.
Mouse embryonic stem cells with aberrant transforming growth factor β signalling exhibit impaired differentiation in vitro and in vivo, 63:101-113

Hayward S.W., Haughney P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.
Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.
Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

Hair follicle

Weiner L., Green H.
Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272
Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Hemidesmosomes

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.
Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Hepatomas

Zvibel I., Fiorino A.S., Brill S., Reid L.M.
Phenotypic characterization of hepatoma cell lines and lineage-specific regulation of gene expression by differentiation agents, 63:213-221

Homeodomains

Bendall A.J., Rincón-Limas D.E., Botas J., Abate-Shen C.
Protein complex formation between *Msx1* and *Lhx2* homeoproteins is incompatible with DNA binding activity, 63:151-157

Human cells

Vrana J.A., Saunders A.M., Chellappan S.P., Grant S.
Divergent effects of bryostatin 1 and phorbol myristate acetate on cell cycle arrest and maturation in human myelomonocytic leukemia cells (U937), 63:33-42

Nohr D., Mai J.K.
Distribution of the CD15 epitope in the mammalian developing lung is opposite in mouse compared with human, 63:43-49

Siggelkow H., Niedhart C., Kurre W., Ihbe A., Schulz A., Atkinson M.J., Hüfner M.
In vitro differentiation potential of a new human osteosarcoma cell line (HOS 58), 63:81-91

Wang Y., Jensen P.J.
Regulation of the level and glycosylation state of plasminogen activator inhibitor type 2 during human keratinocyte differentiation, 63:93-99

Hayward S.W., Haughney P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.
Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Dumortier J., Daemi N., Pourreya C., Anderson W., Bellaton C., Jacquier M.-F., Bertrand S., Chayvialle J.-A., Remy L.
Loss of epithelial differentiation markers and acquisition of vimentin expression after xenograft with laminin-1 enhance migratory and invasive abilities of human colon cancer cells LoVo C5, 63:141-150

Magnaldo T., Fowles D., Darmon M.
Galectin-7, a marker of all types of stratified epithelia, 63:159-168

Stingl J., Eaves C.J., Kuusk U., Emerman J.T.
Phenotypic and functional characterization in vitro of a multipotent epithelial cell present in the normal adult human breast, 63:201-212

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Rognoni J.-B., Pichard V., Honore S., Rigot V., Lehmann M., Roccabianca M., Carles G., Luis J., Marvaldi J., Briand C.
Convergent effects of growth factors, hormones, and fibronectin are necessary for the enterocyte differentiation of a colon adenocarcinoma cell line (HT29-D4), 63:305-317

Hynobius

Ohmura H., Wakahara M.
Transformation of skin from larval to adult types in normally metamorphosing and metamorphosis-arrested salamander, *Hynobius retardatus*, 63:237-246

Insects

Sreng L.

Apoptosis-inducing brain factors in maturation of an insect sex pheromone gland during differentiation, 63:53-58

Intercalated cells

Kloth S., Gmeiner T., Aigner J., Jennings M.L., Röckl W., Minuth W.W.

Transitional stages in the development of the rabbit renal collecting duct, 63:21-32

Intestine

Fabre-Jonca N., Allaman J.-M., Radlgruber G., Meda P., Kiss J.Z., French L.E., Masson D.

The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Krause R., Hemberger M., Messerschmid M., Mayer W., Kothary R., Dixkens C., Fundele R.

Molecular cloning and characterization of murine *Mpgc60*, a gene predominantly expressed in the intestinal tract, 63:285-294

Invasion

Dumortier J., Daemi N., Pourreyaon C., Anderson W., Bellaton C., Jacquier M.-F., Bertrand S., Chayvialle J.-A., Remy L.

Loss of epithelial differentiation markers and acquisition of vimentin expression after xenograft with laminin-1 enhance migratory and invasive abilities of human colon cancer cells LoVo C5, 63:141-150

Kazal type trypsin inhibitor

Krause R., Hemberger M., Messerschmid M., Mayer W., Kothary R., Dixkens C., Fundele R.

Molecular cloning and characterization of murine *Mpgc60*, a gene predominantly expressed in the intestinal tract, 63:285-294

Keratinocytes

Wang Y., Jensen P.J.

Regulation of the level and glycosylation state of plasminogen activator inhibitor type 2 during human keratinocyte differentiation, 63:93-99

Keratins

Appert A., Fridmacher V., Locquet O., Magre S.

Patterns of keratins 8, 18 and 19 during gonadal differentiation in the mouse: sex- and time-dependent expression of keratin 19, 63:273-284

Kidney

Kloth S., Gmeiner T., Aigner J., Jennings M.L., Röckl W., Minuth W.W.

Transitional stages in the development of the rabbit renal collecting duct, 63:21-32

Fabre-Jonca N., Allaman J.-M., Radlgruber G., Meda P., Kiss J.Z., French L.E., Masson D.

The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Laminin integrin receptors

Schünemann H.J., Dillon D., Nielsen L.C., Lwebuga-Mukasa J.S. Modulation of laminin integrin receptors in the postnatal and adult rat lung, 63:181-191

Lens

Mizuno N., Mochii M., Takagi C., Takahashi T.C., Eguchi G., Okada T.S.

A critical role for the optic vesicle in lens development; A reinvestigation of free lens formation in *Cynops pyrrhogaster*, 63:247-252

Limb

Bendall A.J., Rincón-Limas D.E., Botas J., Abate-Shen C.

Protein complex formation between *Msx1* and *Lhx2* homeoproteins is incompatible with DNA binding activity, 63:151-157

LIM-domain

Bendall A.J., Rincón-Limas D.E., Botas J., Abate-Shen C.

Protein complex formation between *Msx1* and *Lhx2* homeoproteins is incompatible with DNA binding activity, 63:151-157

Liver-specific genes

Zvibel I., Fiorino A.S., Brill S., Reid L.M.

Phenotypic characterization of hepatoma cell lines and lineage-specific regulation of gene expression by differentiation agents, 63:213-221

Lung development

Nohr D., Mai J.K.

Distribution of the CD15 epitope in the mammalian developing lung is opposite in mouse compared with human, 63:43-49

Schünemann H.J., Dillon D., Nielsen L.C., Lwebuga-Mukasa J.S.

Modulation of laminin integrin receptors in the postnatal and adult rat lung, 63:181-191

Mammary gland

Stingl J., Eaves C.J., Kuusk U., Emerman J.T.

Phenotypic and functional characterization in vitro of a multipotent epithelial cell present in the normal adult human breast, 63:201-212

Metamorphosis

Ishizuya-Oka A., Inokuchi T., Ueda S.

Thyroid hormone-induced apoptosis of larval cells and differentiation of pepsinogen-producing cells in the stomach of *Xenopus laevis* in vitro, 63:59-68

Ohmura H., Wakahara M.

Transformation of skin from larval to adult types in normally metamorphosing and metamorphosis-arrested salamander, *Hynobius retardatus*, 63:237-246

Microtubule-associated proteins

Fabre-Jonca N., Allaman J.-M., Radlgruber G., Meda P., Kiss J.Z., French L.E., Masson D.

The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Microtubules

Fabre-Jonca N., Allaman J.-M., Radlgruber G., Meda P., Kiss J.Z., French L.E., Masson D.

The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Mid-blastula transition

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.

Post-transcriptional control of *c-myc* RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69-79

Molar development

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.

Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Mouse development

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.

Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Nohr D., Mai J.K.

Distribution of the CD15 epitope in the mammalian developing lung is opposite in mouse compared with human, 63:43-49

Goumans M.-J., Ward-van Oostwaard D., Wianny F., Savatier P., Zwijsen A., Mummery C.

Mouse embryonic stem cells with aberrant transforming growth factor β signalling exhibit impaired differentiation in vitro and in vivo, 63:101-113

Cohen-Tanugi A., Forest N.

Retinoic acid suppresses the osteogenic differentiation capacity of murine osteoblast-like 3A/1D-1M cell cultures, 63:115-123

Fabre-Jonca N., Allaman J.-M., Radlgruber G., Meda P., Kiss J.Z., French L.E., Masson D.

The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.

Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Weiner L., Green H.

Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272

Appert A., Fridmacher V., Locquet O., Magre S.

Patterns of keratins 8, 18 and 19 during gonadal differentiation in the mouse: sex- and time-dependent expression of keratin 19, 63:273-284

Krause R., Hemberger M., Messerschmid M., Mayer W., Kothary R., Dixkens C., Fundele R.

Molecular cloning and characterization of murine *Mpgc60*, a gene predominantly expressed in the intestinal tract, 63:285-294

Mpgc60

Krause R., Hemberger M., Messerschmid M., Mayer W., Kothary R., Dixkens C., Fundele R.

Molecular cloning and characterization of murine *Mpgc60*, a gene predominantly expressed in the intestinal tract, 63:285-294

MyC

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.

Post-transcriptional control of c-myc RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69-79

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.

Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Myoid cells

Galdieri M., Ricci G.

Characterization of different cell populations isolated from rat testis peritubular cells, 63:13-19

Neoteny

Ohmura H., Wakahara M.

Transformation of skin from larval to adult types in normally metamorphosing and metamorphosis-arrested salamander, *Hynobius retardatus*, 63:237-246

Optic vesicle

Mizuno N., Mochii M., Takagi C., Takahashi T.C., Eguchi G., Okada T.S.

A critical role for the optic vesicle in lens development; A reinvestigation of free lens formation in *Cynops pyrrhogaster*, 63:247-252

Organ culture

Ishizuya-Oka A., Inokuchi T., Ueda S.

Thyroid hormone-induced apoptosis of larval cells and differentiation of pepsinogen-producing cells in the stomach of *Xenopus laevis* in vitro, 63:59-68

Osteoblast

Cohen-Tanugi A., Forest N.

Retinoic acid suppresses the osteogenic differentiation capacity of murine osteoblast-like 3A/1D-1M cell cultures, 63:115-123

Osteoma

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.

Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Osteopontin

Siggelkow H., Niedhart C., Kurre W., Ihbe A., Schulz A., Atkinson M.J., Hüfner M.

In vitro differentiation potential of a new human osteosarcoma cell line (HOS 58), 63:81-91

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.

Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Outer hair root sheath

Weiner L., Green H.

Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272

Ovary

Fröjdman K., Pelliniemi L.J., Virtanen I.

Differential distribution of type IV collagen chains in the developing rat testis and ovary, 63:125-130

Appert A., Fridmacher V., Locquet O., Magre S.

Patterns of keratins 8, 18 and 19 during gonadal differentiation in the mouse: sex- and time-dependent expression of keratin 19, 63:273-284

p21^{CIP1/WAF1}

Vrana J.A., Saunders A.M., Chellappan S.P., Grant S.

Divergent effects of bryostatin 1 and phorbol myristate acetate on cell cycle arrest and maturation in human myelomonocytic leukemia cells (U937), 63:33-42

Pepsinogen

Ishizuya-Oka A., Inokuchi T., Ueda S.

Thyroid hormone-induced apoptosis of larval cells and differentiation of pepsinogen-producing cells in the stomach of *Xenopus laevis* in vitro, 63:59-68

Peritubular cells

Galdieri M., Ricci G.

Characterization of different cell populations isolated from rat testis peritubular cells, 63:13-19

Phorbol ester

Vrana J.A., Saunders A.M., Chellappan S.P., Grant S.

Divergent effects of bryostatin 1 and phorbol myristate acetate on cell cycle arrest and maturation in human myelomonocytic leukemia cells (U937), 63:33-42

Pilosebaceous unit

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Plasminogen activator inhibitor 2 (PAI-2)

Wang Y., Jensen P.J.
Regulation of the level and glycosylation state of plasminogen activator inhibitor type 2 during human keratinocyte differentiation, 63:93-99

Pneumocytes type II

Schünemann H.J., Dillon D., Nielsen L.C., Lwebuga-Mukasa J.S.
Modulation of laminin integrin receptors in the postnatal and adult rat lung, 63:181-191

Preadipocytes

Nakajima I., Yamaguchi T., Ozutsumi K., Aso H.
Adipose tissue extracellular matrix: newly organized by adipocytes during differentiation, 63:193-200

Principal cells

Kloth S., Gmeiner T., Aigner J., Jennings M.L., Röckl W., Minuth W.W.
Transitional stages in the development of the rabbit renal collecting duct, 63:21-32

Procollagen I

Siggelkow H., Niedhart C., Kurre W., Ihbe A., Schulz A., Atkinson M.J., Hüfner M.
In vitro differentiation potential of a new human osteosarcoma cell line (HOS 58), 63:81-91

Proliferation control

Vrana J.A., Saunders A.M., Chellappan S.P., Grant S.
Divergent effects of bryostatin 1 and phorbol myristate acetate on cell cycle arrest and maturation in human myelomonocytic leukemia cells (U937), 63:33-42

Promyelocytic cells

Vrana J.A., Saunders A.M., Chellappan S.P., Grant S.
Divergent effects of bryostatin 1 and phorbol myristate acetate on cell cycle arrest and maturation in human myelomonocytic leukemia cells (U937), 63:33-42

Prostate

Hayward S.W., Haughey P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.
Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Protein-protein interactions

Bendall A.J., Rincón-Limas D.E., Botas J., Abate-Shen C.
Protein complex formation between Msx1 and Lhx2 homeoproteins is incompatible with DNA binding activity, 63:151-157

Rabbit development

Kloth S., Gmeiner T., Aigner J., Jennings M.L., Röckl W., Minuth W.W.
Transitional stages in the development of the rabbit renal collecting duct, 63:21-32

Rat development

Fausser J.-L., Schlepp O., Aberdam D., Meneguzzi G., Ruch J.-V., Lesot H.
Localization of antigens associated with adherens junctions, desmosomes, and hemidesmosomes during murine molar morphogenesis, 63:1-11

Galdieri M., Ricci G.

Characterization of different cell populations isolated from rat testis peritubular cells, 63:13-19

Fröjdman K., Pelliniemi L.J., Virtanen I.

Differential distribution of type IV collagen chains in the developing rat testis and ovary, 63:125-130

Hayward S.W., Haughey P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.

Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Schünemann H.J., Dillon D., Nielsen L.C., Lwebuga-Mukasa J.S.
Modulation of laminin integrin receptors in the postnatal and adult rat lung, 63:181-191

Zvibel I., Fiorino A.S., Brill S., Reid L.M.

Phenotypic characterization of hepatoma cell lines and lineage-specific regulation of gene expression by differentiation agents, 63:213-221

Etienne P., Parés-Herbuté N., Mani-Ponset L., Gabrion J., Rabesandratana H., Herbuté S., Monnier L.
Phenotype modulation in primary cultures of aortic smooth muscle cells from streptozotocin-diabetic rats, 63:223-234

Retinoic acid

Cohen-Tanugi A., Forest N.
Retinoic acid suppresses the osteogenic differentiation capacity of murine osteoblast-like 3A/1D-1M cell cultures, 63:115-123

Retrovirus

Goralczyk R., Appold A., Luz A., Riemann S., Strauss P.G., Erfle V., Schmidt J.
Establishment and characterization of osteoblast-like cell lines from retrovirus (RFB MuLV)-induced osteomas in mice, 63:253-262

Serum effect

Galdieri M., Ricci G.
Characterization of different cell populations isolated from rat testis peritubular cells, 63:13-19

Sex differentiation

Fröjdman K., Pelliniemi L.J., Virtanen I.
Differential distribution of type IV collagen chains in the developing rat testis and ovary, 63:125-130

Appert A., Fridmacher V., Locquet O., Magre S.
Patterns of keratins 8, 18 and 19 during gonadal differentiation in the mouse: sex- and time-dependent expression of keratin 19, 63:273-284

Sex pheromones

Sreng L.
Apoptosis-inducing brain factors in maturation of an insect sex pheromone gland during differentiation, 63:53-58

Skeletal muscle

Nakajima I., Yamaguchi T., Ozutsumi K., Aso H.
Adipose tissue extracellular matrix: newly organized by adipocytes during differentiation, 63:193-200

Skin

Ohmura H., Wakahara M.
Transformation of skin from larval to adult types in normally metamorphosing and metamorphosis-arrested salamander, *Hynobius retardatus*, 63:237-246

Kurzen H., Moll I., Moll R., Schäfer S., Simics E., Amagai M., Wheelock M.J., Franke W.W.
Compositionally different desmosomes in the various compartments of the human hair follicle, 63:295-304

Weiner L., Green H.

Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272

Smooth muscle

Hayward S.W., Haughney P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.

Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Etienne P., Parés-Herbuté N., Mani-Ponset L., Gabrion J., Rabesandratana H., Herbuté S., Monnier L.

Phenotype modulation in primary cultures of aortic smooth muscle cells from streptozotocin-diabetic rats, 63:223-234

Stem cells

Stingl J., Eaves C.J., Kuusk U., Emerman J.T.

Phenotypic and functional characterization in vitro of a multipotent epithelial cell present in the normal adult human breast, 63:201-212

Zvibel I., Fiorino A.S., Brill S., Reid L.M.

Phenotypic characterization of hepatoma cell lines and lineage-specific regulation of gene expression by differentiation agents, 63:213-221

Weiner L., Green H.

Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272

Stomach

Ishizuya-Oka A., Inokuchi T., Ueda S.

Thyroid hormone-induced apoptosis of larval cells and differentiation of pepsinogen-producing cells in the stomach of *Xenopus laevis* in vitro, 63:59-68

Stratified epithelia

Magnaldo T., Fowlis D., Darmon M.

Galectin-7, a marker of all types of stratified epithelia, 63:159-168

Weiner L., Green H.

Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272

Stroma

Hayward S.W., Haughney P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.

Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Testis

Galdieri M., Ricci G.

Characterization of different cell populations isolated from rat testis peritubular cells, 63:13-19

Fröjdman K., Pelliniemi L.J., Virtanen I.

Differential distribution of type IV collagen chains in the developing rat testis and ovary, 63:125-130

Fabre-Jonca N., Allaman J.-M., Radgruber G., Meda P., Kiss J.Z., French L.E., Masson D.

The distribution of murine 115-kDa epithelial microtubule-associated protein (E-MAP-115) during embryogenesis and in adult organs suggests a role in epithelial polarization and differentiation, 63:164-180

Appert A., Fridmacher V., Locquet O., Magre S.

Patterns of keratins 8, 18 and 19 during gonadal differentiation in the mouse: sex- and time-dependent expression of keratin 19, 63:273-284

Tissue recombinations

Hayward S.W., Haughney P.C., Rosen M.A., Greulich K.M., Weier H.-U.G., Edahiya R., Cunha G.R.

Interactions between adult human prostatic epithelium and rat urogenital sinus mesenchyme in a tissue recombination model, 63:131-140

Transcription factors

Zvibel I., Fiorino A.S., Brill S., Reid L.M.

Phenotypic characterization of hepatoma cell lines and lineage-specific regulation of gene expression by differentiation agents, 63:213-221

Transforming growth factor β receptor

Goumans M.-J., Ward-van Oostwaard D., Wianny F., Savatier P., Zwijsen A., Mummery C.

Mouse embryonic stem cells with aberrant transforming growth factor β signalling exhibit impaired differentiation in vitro and in vivo, 63:101-113

Vimentin

Dumortier J., Daemi N., Pourreyron C., Anderson W., Bellaton C., Jacquier M.-F., Bertrand S., Chayvialle J.-A., Remy L.

Loss of epithelial differentiation markers and acquisition of vimentin expression after xenograft with laminin-1 enhance migratory and invasive abilities of human colon cancer cells LoVo C5, 63:141-150

Xenopus

Ishizuya-Oka A., Inokuchi T., Ueda S.

Thyroid hormone-induced apoptosis of larval cells and differentiation of pepsinogen-producing cells in the stomach of *Xenopus laevis* in vitro, 63:59-68

Andéol Y., Lefresne J., Simard C., Séguin C., Mouton C., Signoret J.

Post-transcriptional control of c-myc RNA during early development analyzed in vivo with *Xenopus*-axolotl heterologous system, 63:69-79

Yolk sac

Goumans M.-J., Ward-van Oostwaard D., Wianny F., Savatier P., Zwijsen A., Mummery C.

Mouse embryonic stem cells with aberrant transforming growth factor β signalling exhibit impaired differentiation in vitro and in vivo, 63:101-113

Zinc-finger protein

Weiner L., Green H.

Basonuclin as a cell marker in the formation and cycling of the murine hair follicle, 63:263-272

